

## Cosmic ray astrophysics

Cosmic rays (CRs) below  $10^{15}$  eV are thought to be Fermi accelerated in supernova remnant shocks, but the origin of CRs above that and up to the ‘GZK’ limit  $E_{siml}10^{20}$  eV is debated. A steepening of the diffuse spectrum attributed to the GZK process (interaction with CMB photons) has been observed by the Auger observatory [2], so the sources appear to be inside  $R_{GZK} \lesssim 50 - 100$ Mpc, and a spatial correlation analysis of arrival directions [1] indicated a weak correlation with certain AGNs and a much firmer correlation with galaxy large scale structures inside that radius. These observations strongly favor an astrophysical, as opposed to a ‘new physics’, origin and the question is what are these sources.

**AGNs**, in particular luminous radio-loud (jetted) AGNs have long been considered candidates [3], but there are only a few of these inside  $R_{GZK}$ . A more promising possibility is that acceleration occurs in **radio-quiet AGNs**, [4], which may provide acceleration sites with the right properties in large enough numbers.

**GRB** are the other main candidates [5, 6, 7]. These are transient, and taking into account diffusion time delays, they occur at a sufficient rate inside  $R_{GZK}$  [8]. The maximum energy of protons achievable, both in AGNs and GRBs, is the GZK value

$$E_p \lesssim \beta_s Z e B R \sim 10^{20.5} \text{ eV}. \quad (1)$$

In GRB the number spectrum expected has slope -2, which is suitable for explaining the diffuse CR flux in the  $10^{19} - 10^{21}$  eV range. Proof, however, is difficult. Neutrinos from  $p\gamma$  at these energies would have  $E_\nu \sim 10^{18}$  eV, which requires larger instruments than, e.g., ANITA. Secondary photon emission from  $p\gamma$  cascades, however, may appear at GeV energies, providing potential CR signatures [15], and Fermi observations of GRB 090510 at GeV energies are indeed compatible with such hadronic GRB models [10]. If GRB indeed accelerate the observed GZK CRs, the supernova remnants of their progenitors could be sites of the late decay of neutrons (produced originally by  $p\gamma$  interactions) back into protons and positrons, the latter leading to synchrotron-IC photon cascades which could be detectable with GeV and/or TeV telescopes [11].

At lower energies the -2 spectrum is flatter than the observed -2.7, and either nearby GRB [9] or other sources must be invoked. The latter may be **hypernovae**, supernovae with semi-relativistic ejecta, which can accelerate protons up to  $10^{19}$  eV or heavy nuclei up to  $10^{20.5}$  eV [12, 14, 13]. The numbers and the inferred distribution of ejecta velocities yield the appropriate spectrum and diffuse flux.

## References

- [1] The Pierre Auger Collaboration et al. 2007, Science, 318:938
- [2] Abraham, J. and the Pierre Auger Coll., 2008, PRL, 101:061101
- [3] Berezhinsky, V, Gazizov, A, and Grigorieva, S, 2006, PRD 74:043005
- [4] Pe’er, A., Murase, K., and Mészáros, P., 2010, PRD, in press (arXiv:0911.1776)
- [5] Waxman, E, 1995, PRL, 75:386
- [6] Vietri, M, 1995, ApJ, 453:883
- [7] Dermer, C, 2002, ApJ 574:65 (astro-ph/0005440)
- [8] Waxman, E., 2006, AIPC 836:589
- [9] Wick, S, Dermer, C, Atoyan, A, 2004, Astropart.Phys. 21:125
- [10] Asano, K; Guiriec, S; Mészáros, P., 2009, ApJ(Lett.), 705:L19
- [11] Ioka, K., Kobayashi, S and Mészáros, P., 2004, ApJ, 613:L171
- [12] Wang, X-Y, Razzaque, S, Mészáros, P and Dai, Z-G, 2007, PRD, 76:083009
- [13] Wang, X-Y, Razzaque, S and Mészáros, P, 2008, ApJ, 677:432
- [14] Budnik, R; Katz, B; MacFadyen, A; Waxman, E, 2008, ApJ 673:928
- [15] Asano, K., Inoue, S. and Mészáros, P., 2009, ApJ, 699:953